

- Anderson, C. I. H. S., G.W. (1998). "The occurrence of distinct morphotypes within a population of *Fucus spiralis*" *Journal of the Marine Biological Association of the UK* 79: 1-4.
- Astbury, W. T. F., R.D. (1948). "The structure of the Cell Wall in Some Species of the Filamentous Green Alga *Cladophora*" *Proceedings of the Royal Society of London. Series B, Biological Sciences* 129(854): 54-76
- Atkins, S. M. (1985). The structure and regulation of infaunal communities off the Yorkshire Coast, Leeds. PhD.
- Barne, J. H., Rabson, C.F., Kazanekow, J.P., Doady, J.P., Davidson, N.C. (eds.) (1995). Coasts and Seas of the United Kingdom. Region 5, north-east England, Berwick upon Tweed to Fliley Bay, Joint Nature Conservation Committee
- Barne, J. H., Rabson, C.F., Kazanekow, J.P., Doady, J.P., Davidson, N.C. (eds.) (1995). Coasts and Seas of the United Kingdom. Region 6 Eastern England: Flamborough Head to Great Yarmouth. Peterborough Joint Nature Conservation Committee
- Bayliss, R. (1995). Draft Technical Report. Flamborough Head Sensitive Marine Area.
- Bennett, T. L. (1991). Benthic Marine ecosystems in Great Britain: A review of present knowledge, Orkney, North Scotland and North East England. Marine Nature Conservation Review. JNC. Peterborough, English Nature Section 2 to 5.
- Bennett, T. L. F.-S., J.L. (1993). South-east Scotland and North-east England (Dunbar to Bridlington) (MNC sector 5). Marine Nature Conservation Review. Benthic Marine ecosystems of Great Britain and the north-east Atlantic. K. Hiscok. Peterborough, Joint Nature Conservation Committee (Coasts and Seas of the United Kingdom. MNC series): 155-177.
- Bird, G. M., E. (1992). Report on the 1991 intertidal invertebrate survey of the EC designated beaches in the Yorkshire Region of the NRA., National Rivers Authority, Yorkshire Region (Now Environment Agency).
- Bowman, R. S. L. J. R. (1986). "Geographical variation in the breeding cycles and recruitment of *Patella* species, in: Long term changes in coastal benthic communities. Cost 647 project on coastal benthic ecology ed. by C.Help, B.F. Keegan, & J.R. Lewis" *Hydrobiologia* 142: 41-56.
- Brown, A. R. C. N. C. D. (1993). Marine Monitoring for Cleveland Potash Ltd.; in 1993 ICI Group Environmental Laboratory, Brixham.
- Clayson, C. H. E. J. A. (1990). Directional Wave Data Recorded off Flamborough Head, Institute of Oceanographic Sciences.
- Covey, R. (1991). Benthic marine ecosystems in Great Britain: a review of current knowledge, eastern England and eastern Channel (MNC coastal sectors 6 and 7) CSD report Marine Nature Conservation Review, Nature Conservancy Council (Now English Nature).
- Craig, M. C. D., Shillabeer, N. & Parr, J.C.W. (1993). "Long term benthic monitoring of the effects of a dense effluent discharged to North Sea coastal waters" *Journal of the Institute of Water and Environmental Management* 7: 262-269.
- Craig, M. C. D. S., N. (1998). Marine Monitoring for Cleveland Potash Ltd., Brixham, ICI Group Environmental Laboratory, Brixham.
- Crimp, T. J. (1985). The ecology of rocky shore marine nematodes. Leeds, University of Leeds. PhD.
- Croagh, F., Johnson L.J., Morris, E. & Scott, G.W. (1994). "Evidence for increased levels of imposex in *Nucella lapillus* (L.) populations on the north-east coast of England." *Naturalist* 119.
- Davies, J. & Sotherton T. (1995). Mapping the distribution of benthic biotopes on Flamborough Head. English Nature Research Report, English Nature.
- Denness, B. (1991). Fliley Bay Wave Regime, Bureau of Applied Sciences.
- Dosser, J. (1966). Report on Proposed Coast protection order: Fliley Bay, Bridlington Rural District Council.
- Edwards, E. (1967). Yorkshire crab stocks. M. o. A. F. a. F. n. (CEFA), Ministry of Agriculture Food and Fisheries, Burnham on Crouch.
- Elliott, M., Jones N.W., Lewis D.S., Pethick J.S., Symes, G. (1991). Fliley Bay Environmental Statement Institute of Estuarine & Coastal Studies, University of Hull.
- Feare, C. J. (1969). The dynamics of a population of dogwhelks (*Thais lapillus* [Now *Nucella lapillus*]), University of Leeds. PhD.
- Feare, C. J. (1970). "Aspects of the ecology of an exposed shore population of dogwhelks *Nucella lapillus* (L)." *Oecologia* (Berlin) 5: 1-18.
- Feare, C. J. (1971). "The adaptive significance of aggregation behaviour in the dogwhelk *Nucella lapillus* (L)." *Oecologia* (Berlin) 7: 117-126.
- Foster-Smith, R. I. (1988). Marine Interpretation on the North York Moors National park coastline, Nature Conservancy Council.
- George, J. D., Frinham, A.A., Tittley, I., & Price, J.H. (1988). The macrobenthos of chalk shores in North Norfolk and around Flamborough Head., British Museum (Natural History) Nature Conservancy Council report (Now English Nature)
- Grahame, J., Hull, S.L., Mill, P.J. & Hastings, R. (1998). Realising unrecognised biodiversity in marine molluscs London Cambridge University Press.
- Grahame, J. H. F. (1985). "Factors affecting the distribution of the epiphytic fauna of *Corallina officinalis* (L) on an exposed rocky shore" *Ophelia* 30: 113-129.
- Gray, J. S. R., R.W. (1971). "A quantitative study of the meiofauna of an exposed sandy beach at Robin Hood's Bay, Yorkshire." *Journal of the Marine Biological Association of the UK* 51: 1-19.
- Hama, P. S. (1989). Studies on the biology of pool dwelling serpulids & spirorbids (*Annulid:polychaeta*) at Fliley Brigg, North Yorkshire. Leeds, University of Leeds. PhD.
- Hardy, F. G. S., G.W. (1994). "Seaweeds of the Yorkshire Coast." *The Phycologist* 38: 22-25.
- Harris, C. (1966). Fliley Bay Erosion Study, University of Hull.
- Hay, W. C. (1894). Conchology: marine land and freshwater shells.
- Hawson, C., Mercer, T. & Burker, F. (2003). Flamborough Head Sublittoral Monitoring, English Nature: 120.
- Hawson, C. M. (2000). Flamborough Head cSAC sea cave survey. English Nature Research Report. Peterborough, English Nature
- Hawson, C. M. (2002). Flamborough Head cSAC Intertidal survey., English Nature Research Report. Peterborough, English Nature.
- Hull, S. L., Grahame, J. & Mill, P.J. (1996). "Morphological divergence and evidence for reproductive isolation in littorine saxatilis (Oliv.) in north-east England" *Journal of Molluscan studies* 62: 89-99.
- Hull, S. L. (1997). "Seasonal changes of diversity and abundance of ostracods on four species of intertidal algae with differing structural complexity." *MARINE ECOLOGY-PROGRESS SERIES* 161: 71-82
- Hull, S. L. (1998). "Assortative mating between two distinct micro-allopatric populations of littorine saxatilis (Oliv.) on the north-east coast of England." *Hydrobiologia* 378: 79-88.
- Hull, S. L. (1999). "Comparison of tidepool physical ostracod abundance and assemblage structure on three spatial scales" *MARINE ECOLOGY-PROGRESS SERIES* 182: 201-208.
- Hull, S. L. (1999). "Intertidal ostracod (Crustacea: Ostracoda) abundance and assemblage structure within and between four shores in north-east England." *Journal of the Marine Biological Association of the UK* 79: 1845-1851.
- Hull, S. L., Grahame, J. & Mill, P.J. (1999). "Reproduction in four populations of brooding pertusella (Littorina) at Ravenscar, North Yorkshire: adaptation to the local environment?" *Journal of the Marine Biological Association of the UK* 79: 891-898.
- Hull, S. L., Winter, L.J., & Scott, G.W. (2001). "Habitat heterogeneity, body size and phenotypic diversity in *Idotea granulosa* (Isopoda) on the north-east coast of England." *Journal of the Marine Biological Association of the UK* 81: 949-954.
- Hull, S. L. R., D. (2000). "Clonal diversity and rockpool size in the marine ostracod, *Gallistocythere baltica*." *Journal of the Marine Biological Association of the UK* 80(3).
- Hull, S. L. R., D. (2000). "Sex-biased colour polymorphism in the marine ostracod *Paradoxostoma varicella* (Crustacea)." *Journal of the Marine Biological Association of the UK* 80: 69-73.
- Hurst, C. D. S., D.O.F. (1995). "Comparison of allozyme and mitochondrial DNA spatial differentiation in the limpet *Patella vulgata*." *Marine Biology* 122(2): 257-263.
- Johnson, L. J. (1999). The population dynamics and ecology of barnacle-dwelling littorinids. University of Hull, Scarborough Scarborough University of Hull
- Kendall, M. A., Bowen, A.S., Williamson, P., & Lewis, J.R. (1985). "Annual variation in recruitment of Semibalanus balanoides on the North Yorkshire Coast, 1969-1981." *Journal of the Marine Biological Association of the United Kingdom. Plymouth* 65(4): 1009-1030.
- Lewis, D. B., et al. (1987). Fliley Brigg, its plants and animals. University of Hull Adult Education department
- Lewis, D. B. (1987). Scarborough Shoreline, University of Hull Department of Adult and continuing education
- Lewis, J. R. (1964). The ecology of rocky shores. London, English University Press.
- Lewis, J. R. (1977). The role of physical and biological factors in the distribution and stability of rocky shore communities 11th European Symposium on Marine Biology, Gtway Pergamon Press Oxford.
- Lockwood, S. (1972). An ecological study of a  $\theta$ -group plaice population, Fliley Bay, Yorkshire, University of East Anglia. PhD.
- Lockwood, S. J. (1975). "The settlement, distribution and movements of  $\theta$ -group plaice *Pleuronectes platessa* in Fliley Bay, Yorkshire." *J. Fish. Biol.* 6: 465-477.
- Masse, G. (1885). "Marine Algae of the Scarborough District." *Naturalist* 10: 300-306.
- McCrorty, S. (1972). Factors affecting the distribution of *Bathyporeia* species. Leeds, University of Leeds. PhD.
- Moore, P. G. (1972). "Particulate matters in the sublittoral zone of an exposed coast and its ecological significance with reference to the fauna inhabiting kelp holdfasts." *J. Exp. Mar. Biol. & Ecol* 18: 59-88.
- Moore, P. G. (1973). "The kelp fauna of north-east Britain II *Multivertebrae* classification: turbidity as an ecological factor." *J. Exp. Mar. Biol. & Ecol* 13: 127-163.
- Moore, P. G. (1973). "The kelp fauna of north-east Britain I: Introduction and the physical environment." *J. Exp. Mar. Biol. & Ecol* 13: 127-163.
- Moore, P. G. (1974). "The kelp fauna of north-east Britain III. Qualitative and Quantitative ordinations and the utility of a multivariate approach." *J. Exp. Mar. Biol. & Ecol* 16: 257-300.
- National Rivers Authority, Y. R. (1980). A biological survey of the inter-tidal zone at Scarborough in relation to existing sewage discharges and the proposed new long sea outfall at Scalby Ness, June - July 1980, National Rivers Authority (Now Environment Agency).
- National Rivers Authority, Y. R. (1991). A biological survey of the inter-tidal zone at Scarborough in relation to existing sewage discharges and the proposed new long sea outfall at Scalby Ness, June - July 1989, National Rivers Authority (Now Environment Agency).
- National Rivers Authority, Y. R. (1991). Levels of TBT induced imposex in the dogwhelk *Nucella lapillus* along the North Yorkshire Coast National Rivers Authority, Yorkshire Region (Now Environment Agency).
- Newton, A. J. (1972). Marine Biological Investigation concerning a proposed industrial effluent, University of Leeds. PhD thesis.
- Norris, A. (1973). "Reports of field trips to Scarborough, South Bay; Thorneick Bay, Flamborough and Fliley Brigg, 1972 Conchological Section marine survey." *The Naturalist*(April/June): 71-73.
- Pearshall, W. H. M., F.A. (1923). "Yorkshire Naturalists at Bridlington." *Naturalist* 797: 205-212.
- Perkins, D. (1993). Marine Algae - Seaweeds The Natural History of the Scarborough District. G. B. W. F. C. Ristington: 67-88.
- Phillip, G. (1994). "Plant associations of the shore near Flamborough." *Naturalist*: Supplement 384: 15-16.
- Podsigan, B. U., R.F. (1974). "Quantitative changes in blood sugar levels of *Crangon vulgaris*" *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology* 93(1): 1-6.
- Rees, H. L. (1975). Unpublished data on the Benthos of Fliley Bay.
- Rees, H. L. (1979). Application of benthic community studies to pollution monitoring off the north-east coast of England London, University of London. PhD.
- Scott, G. W., MR; Seaneey, B (1990). "Avian habituation to recreational disturbance on the North Yorkshire coast" *Naturalist* 121(1816): 11-15.
- Scott, G. W., Hull, S.L., Hornby, S.E., Gavin Hardy, F. & Deans, N.J.P. (2001). "Phenotypic variation in *Fucus spiralis* (Phaeophyceae): morphology, chemical phenotype and their relationship to the environment." *European Journal of Phycology* 36: 43-58.
- Scott, G. W. H., F.G. (1994). "Observations on the occurrence of hybrids between two sympatric species of *Fucoid* algae." *Cryptogamiae Algologica* 15(4): 297-305.
- Seed, R. (1967). The ecology of *Mytilus edulis* (L) on exposed shores University of Leeds. PhD.
- Seed, R. (1969). "The ecology of *Mytilus edulis* (L) (Lamellibranchiata) on exposed rock shores: II growth and mortality." *Oecologia* (Berlin) 3: 317-350.
- Shillabeer, N. C., M.C.D. (1992). Marine monitoring for Cleveland Potash Ltd.; in 1991. Brixham, ICI Group Environmental Laboratory, Brixham.
- Tittley, I. (1988). Chalk cliff algal communities: 2 Outside south-eastern England. CSD Report. London, British Museum (Natural History) department of Botany.
- Todd, C. D. (1977). The ecology of inter tidal mudflats. Leeds, University of Leeds. PhD.
- Walsh, G. B. R., F.C. (1956). The Natural History of the Scarborough District - Volume 2. .
- Willows, R. I. (1984). The population biology of *Ligia oceanica* (L), University of Leeds. PhD.
- Wisey Environmental (1993). Benthic Survey data, Scarborough 1989 and 1990 Yorkshire Water
- Wood, E. (1988). Flamborough Head sub-littoral survey. CSD report, Nature Conservancy Council (English Nature).
- Woods, F. H. (1912). "The Marine Mollusca of the Yorkshire Coast - as represented in the Hull Museum." *Hull, Scientific and Field Naturalists Club*: 231-249.